

# Convex set


Выпуклые множества.

## Line segment

Выпуклые функции

Suppose  $x_1, x_2$  are two points in  $\mathbb{R}^n$ . Then the line segment between them is defined as follows:

$$x = \theta x_1 + (1 - \theta)x_2, \theta \in [0, 1]$$



# Convex set

The set  $S$  is called **convex** if for any  $x_1, x_2$  from  $S$  the line segment between them also lies in  $S$ , i.e.


Мн-во  $S$  - выпукло тогда и только тогда, когда

$$\forall \theta \in [0, 1], \forall x_1, x_2 \in S: \theta x_1 + (1 - \theta)x_2 \in S$$

$\forall$  отрезок мн-ва  $S$   
 отрезок между любыми  
 элементами лежит в  $S$

## Examples:

- Any affine set
- Ray
- Line segment




$\emptyset$   
 Выпукло по опр




1.



2.




3.




4.



5.




6.



BRO




NOT BRO




BRO



NOT BRO



BRO



BRO

## Related definitions

### Convex combination

Let  $x_1, x_2, \dots, x_k \in S$ , then the point  $\theta_1 x_1 + \theta_2 x_2 + \dots + \theta_k x_k$  is called the convex combination of points  $x_1, x_2, \dots, x_k$  if  $\sum_{i=1}^k \theta_i = 1$ ,  $\theta_i \geq 0$


### Convex hull

The set of all convex combinations of points from  $S$  is called the convex hull of the set  $S$ .


$$\mathbf{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \sum_{i=1}^k \theta_i = 1, \theta_i \geq 0 \right\}$$

- The set  $\mathbf{conv}(S)$  is the smallest convex set containing  $S$ .
- The set  $S$  is convex if and only if  $S = \mathbf{conv}(S)$ .


Examples:




BRO




BRO




BRO



BRO



BRO



BRO

## Finding convexity

In practice it is very important to understand whether a specific set is convex or not. Two approaches are used for this depending on the context.

- By definition.
- Show that  $S$  is derived from simple convex sets using operations that preserve convexity.

### 1) By definition

$$x_1, x_2 \in S, 0 \leq \theta \leq 1 \rightarrow \theta x_1 + (1 - \theta)x_2 \in S$$

## Preserving convexity

### 3) The linear combination of convex sets is convex

Let there be 2 convex sets  $S_x, S_y$ , let the set  $S = \{s \mid s = c_1x + c_2y, x \in S_x, y \in S_y, c_1, c_2 \in \mathbb{R}\}$

Take two points from  $S$ :  $s_1 = c_1x_1 + c_2y_1, s_2 = c_1x_2 + c_2y_2$  and prove that the segment between them  $\theta s_1 + (1 - \theta)s_2, \theta \in [0,1]$  also belongs to  $S$

$$\theta s_1 + (1 - \theta)s_2$$

$$\theta(c_1x_1 + c_2y_1) + (1 - \theta)(c_1x_2 + c_2y_2)$$

$$c_1(\theta x_1 + (1 - \theta)x_2) + c_2(\theta y_1 + (1 - \theta)y_2)$$

$$c_1x + c_2y \in S$$

### 2) The intersection of any (!) number of convex sets is convex

If the desired intersection is empty or contains one point, the property is proved by definition. Otherwise, take 2 points and a segment between them. These points must lie in all intersecting sets, and since they are all convex, the segment between them lies in all sets and, therefore, in their intersection.

## The image of the convex set under affine mapping is convex

$$S \subseteq \mathbb{R}^n \text{ convex} \rightarrow f(S) = \{f(x) \mid x \in S\} \text{ convex} \quad (f(x) = \mathbf{A}x + \mathbf{b})$$

Examples of affine functions: extension, projection, transposition, set of solutions of linear matrix inequality  $\{x \mid x_1 A_1 + \dots + x_m A_m \preceq B\}$  Here  $A_i, B \in \mathbb{S}^p$  are symmetric matrices  $p \times p$ .

Note also that the prototype of the convex set under affine mapping is also convex.

$$S \subseteq \mathbb{R}^m \text{ convex} \rightarrow f^{-1}(S) = \{x \in \mathbb{R}^n \mid f(x) \in S\} \text{ convex} \quad (f(x) = \mathbf{A}x + \mathbf{b})$$

### Example 1

Prove, that ball in  $\mathbb{R}^n$  (i.e. the following set  $\{x \mid \|x - x_c\| \leq r\}$ ) - is convex.

Решение:

1) Возьмём  $x_1 \in S$   $\|x_1 - x_c\| \leq r$   
 $x_2 \in S$   $\|x_2 - x_c\| \leq r$

2) Построим отрезок:  $x = \theta x_1 + (1-\theta)x_2$   $0 \leq \theta \leq 1$

3) Проверим, что  $x \in S$   $\|x - x_c\| \leq r$   $\theta + 1 - \theta = 1$

$$\begin{aligned} \|\theta x_1 + (1-\theta)x_2 - x_c\| &\leq r \rightarrow \|\theta x_1 + (1-\theta)x_2 - \theta x_c - (1-\theta)x_c\| = \\ &= \|\theta(x_1 - x_c) + (1-\theta)(x_2 - x_c)\| \leq \theta \|x_1 - x_c\| + (1-\theta) \|x_2 - x_c\| = \\ &\leq \theta r + (1-\theta)r \leq r \rightarrow x \in S \end{aligned}$$

$\forall \theta!$

### Example 2

$\Rightarrow$  НИ-ВО ВЫПЯТНО!

Which of the sets are convex: 1. Stripe,  $\{x \in \mathbb{R}^n \mid \alpha \leq a^\top x \leq \beta\}$  1. Rectangle,

$\{x \in \mathbb{R}^n \mid \alpha_i \leq x_i \leq \beta_i, i = \overline{1, n}\}$  1. Kleen,  $\{x \in \mathbb{R}^n \mid a_1^\top x \leq b_1, a_2^\top x \leq b_2\}$  1. A set of points

closer to a given point than a given set that does not contain a point,

$\{x \in \mathbb{R}^n \mid \|x - x_0\|_2 \leq \|x - y\|_2, \forall y \in S \subseteq \mathbb{R}^n\}$  1. A set of points, which are closer to one set

than another,  $\{x \in \mathbb{R}^n \mid \text{dist}(x, S) \leq \text{dist}(x, T), S, T \subseteq \mathbb{R}^n\}$  1. A set of points,

$\{x \in \mathbb{R}^n \mid x + X \subseteq S\}$ , where  $S \subseteq \mathbb{R}^n$  is convex and  $X \subseteq \mathbb{R}^n$  is arbitrary. 1. A set of points whose

distance to a given point does not exceed a certain part of the distance to another given point is

$\{x \in \mathbb{R}^n \mid \|x - a\|_2 \leq \theta \|x - b\|_2, a, b \in \mathbb{R}^n, 0 \leq \theta \leq 1\}$

Множество решений системы лнн. ур-ий:

$$S = \{x \in \mathbb{R}^n \mid Ax = b\} \quad S - \text{выпукло?}$$

$$A \in \mathbb{R}^{m \times n} \\ m < n$$

$$1. \quad x_1 \in S \quad Ax_1 = b \\ x_2 \in S \quad Ax_2 = b$$

$$2. \quad x = \theta x_1 + (1-\theta)x_2 \quad 0 \leq \theta \leq 1$$

$$x \in S \quad Ax = b$$

$$A(\theta x_1 + (1-\theta)x_2) \stackrel{?}{=} b$$

$$\theta Ax_1 + (1-\theta)Ax_2 \stackrel{?}{=} b$$

$$\theta b + (1-\theta)b \stackrel{?}{=} b$$

$$b = b$$

Мн-во симм. пол. <sup>полу</sup>отр. матрицы

$$S = \{X \in \mathbb{R}^{n \times n} \mid \forall y \in \mathbb{R}^n: y^T X y \geq 0\}$$

$S$  выпукло!

$$1. \quad X_1 \in S$$

$$y_1^T X_1 y_1 \geq 0 \quad \forall y_1 = y$$

$$2. \quad X_2 \in S$$

$$y_2^T X_2 y_2 \geq 0 \quad \forall y_2 = y$$

$$3. \quad X = \theta X_1 + (1-\theta)X_2 \quad \forall y = y_1 = y_2: y^T X y \geq 0?$$

$$y^T [\theta X_1 + (1-\theta)X_2] y = \theta \underbrace{y^T X_1 y}_{\geq 0} + (1-\theta) \underbrace{y^T X_2 y}_{\geq 0} \geq 0$$

Вероятностный симплекс

Выпукло!


$$S = \{p \in \mathbb{R}^n \mid p \geq 0, \mathbf{1}^T p = 1\}$$

A. доказать выпуклость

$$S_A = \{p \in \mathbb{R}^n \mid p \geq 0\}$$

$$\begin{array}{l} \perp \\ p_i \geq 0 \\ \forall i=1, n \end{array}$$

$$\sum_{i=1}^n p_i = 1$$



B. доказать выпуклость

$$S_B = \{p \in \mathbb{R}^n \mid \mathbf{1}^T p = 1\}$$

$$0 \leq \theta \leq 1$$

$$\mathbf{1}^T p_1 = 1$$

$$\mathbf{1}^T p_2 = 1$$

$$p = \theta p_1 + (1-\theta) p_2$$

$$\mathbf{1}^T p \stackrel{?}{=} 1 \quad p \in S_B$$

$$\begin{aligned} \mathbf{1}^T (\theta p_1 + (1-\theta) p_2) &= \\ &= \theta \mathbf{1}^T p_1 + (1-\theta) \mathbf{1}^T p_2 = \end{aligned}$$

$$= \theta \cdot 1 + (1-\theta) \cdot 1 = \mathbf{1}$$